

A BCS CONDENSATE IN NJL₃₊₁?

SIMON HANDS* AND DAVID N. WALTERS

*Department of Physics, University of Wales Swansea,
Singleton Park, Swansea SA2 8PP, U.K.*

We present results from a lattice Monte Carlo study of the Nambu – Jona-Lasinio model in 3+1 dimensions with a baryon chemical potential $\mu \neq 0$. As μ is increased there is a transition from a chirally-broken phase to relativistic quark matter, in which baryon number symmetry appears spontaneously broken by a diquark condensate at the Fermi surface, implying a superfluid ground state. Finite volume corrections to this relativistic BCS scenario, however, are anomalously large.

The proposal that at high baryon density a condensation of diquark pairs in degenerate quark matter takes place, leading to color superconducting ground states, has generated a profusion of theoretical and phenomenological interest¹. Most approaches assume the existence of a Fermi surface and then consider its instability with respect to diquark condensation arising from an attractive qq interaction. The resulting excitation energy gap Δ can be calculated self-consistently in close analogy to the BCS gap in electronic superconductors. Model calculations yield gaps as large as 50 - 100MeV².

A first-principles calculation of the ground state of nuclear or quark matter using the methods of lattice QCD, however, is hampered by the “sign problem”, ie. $\exp(-S/\hbar)$ is no longer positive definite, and cannot be used as a probability measure in importance sampling the Euclidean path integral, once baryon chemical potential $\mu \neq 0$. It is still possible, however, to study simpler models of strong interactions with $\mu \neq 0$ using lattice methods³. The Nambu – Jona-Lasino (NJL) model, which closely resembles the original BCS model, is particularly interesting in the current context, because it is the only simulable model which exhibits a Fermi

*supported by a PPARC Senior Research Fellowship

surface for sufficiently large μ . The Lagrangian density is

$$L = \bar{\psi}(\not{\partial} + m_0 + \mu\gamma_0)\psi - \frac{g^2}{2} [(\bar{\psi}\psi)^2 - (\bar{\psi}\gamma_5\vec{\tau}\psi)^2]. \quad (1)$$

Here ψ and $\bar{\psi}$ are isospinor “quark” fields carrying baryon charge; note that both $q\bar{q}$ and qq interactions are attractive in the isoscalar channel.

For coupling strength g^2 sufficiently large the vacuum ground state has chiral $SU(2)_L \otimes SU(2)_R$ symmetry spontaneously broken to $SU(2)_{isospin}$ by generation of a chiral condensate and resulting “constituent” quark mass $\Sigma \simeq g^2 \langle \bar{\psi}\psi \rangle$. At zero temperature this state is stable as μ is increased until at $\mu_c \sim \Sigma$ chiral symmetry is restored, at which point the baryon density $n_B = \langle \bar{\psi}\gamma_0\psi \rangle$ rises from zero⁴. For $\mu > \mu_c$ the ground state is thus relativistic quark matter with Fermi momentum $k_F \sim \mu$. A natural question is whether qq attraction in this phase leads to a diquark condensate spontaneously breaking the $U(1)_B$ symmetry of (1) leading (since in this case the broken symmetry is global rather than local) to superfluidity.

The NJL model in 2+1 dimensions has been studied extensively using Monte Carlo simulations. While there is enhanced qq pairing in the scalar isoscalar channel for $\mu > \mu_c$ ⁵, simulations with an explicit $U(1)_B$ -violating source term jqq found no evidence for diquark condensation in the $j \rightarrow 0$ limit⁶. Instead, the condensate scales approximately as $\langle qq(j) \rangle \propto j^{\frac{1}{\delta}}$, the exponent δ increasing with μ . Moreover, the quasiparticle dispersion relation in the spin- $\frac{1}{2}$ channel is consistent with a vanishing BCS gap $\Delta = 0$. The interpretation is that long-wavelength fluctuations of the complex phase of the qq wavefunction destroy the condensate, but that long-range phase coherence remains (ie. correlations are dominated by “spin-waves” and remain algebraic functions of spatial separation) just as in the low- T phase of the 2d XY model. Superfluidity in two dimensions is realised in Kosterlitz-Thouless mode; namely, persistent flow patterns can only be disrupted by creating a vortex/anti-vortex excitation and then translating one member of the pair around the universe, which in the low- T phase entails an energy cost increasing logarithmically with system size.

This rather exotic scenario, namely gapless thin film superfluidity, is inherently non-perturbative and is not exposed by self-consistent approaches. It thus justifies the application of numerical simulations, and begs the question of whether similar surprises will emerge from a study of the phenomenologically-motivated case of NJL_{3+1} ⁷. In 3+1d the model is non-renormalisable, and hence its physical predictions are sensitive to the details of the ultra-violet cutoff. We have matched our lattice model to

low-energy QCD following the procedure of Klevansky ⁴, working to leading order in an expansion in $1/N_c$, where N_c is the number of quark “colors”, and calculating Σ , m_π and f_π in terms of bare parameters g^2 and m_0 . Since we use staggered lattice fermions $\chi, \bar{\chi}$, we have $N_c = 4$, in which case reasonable low-energy phenomenology emerges for the choice $a^2/g^2 = 0.565$, $m_0 a = 0.002$, with inverse lattice spacing $a^{-1} \simeq 1\text{GeV}$. Figure 1 shows




Figure 1. Chiral condensate $\langle \bar{\chi}\chi \rangle$, baryon density, and diquark condensate as functions of μ , showing both large- N_c solution (solid curve) and simulation results (points).

results for $\langle \bar{\chi}\chi \rangle$ and n_B and confirms the transition between vacuum and quark matter at $\mu_c \sim \Sigma \sim \frac{g^2}{2} \langle \bar{\chi}\chi \rangle$. The discrepancy between line and points is ascribed to $O(1/N_c)$ corrections and amounts to a 30% effect. In contrast to $2+1d$ where a sharp first-order transition is observed ^{3,5}, the transition appears continuous in the chiral limit $m_0 \rightarrow 0$, although this may be sensitive to the cutoff ⁴.

To investigate the diquark sector, we introduce a source term $j_\pm(q^{tr}q \pm \bar{q}\bar{q}^{tr})$ into the quark operator used for measurements (since the simulation dynamics are performed with $j_\pm = 0$, this is a “partially quenched” approximation) and define the following:

$$\langle qq_\pm \rangle = \frac{1}{V} \frac{\partial \ln Z}{\partial j_\pm}; \quad \chi_\pm = \frac{\partial \langle qq_\pm \rangle}{\partial j_\pm}; \quad R(j_+) = \left| \frac{\chi_+}{\chi_-} \right|_{j_-=0}. \quad (2)$$

The nature of the ground state is then determined by

$$\begin{aligned} \lim_{j_+ \rightarrow 0} \langle qq_+ \rangle &= 0, \quad \lim_{j_+ \rightarrow 0} R = 1 && \text{if } U(1)_B \text{ manifest} \\ \lim_{j_+ \rightarrow 0} \langle qq_+ \rangle &\neq 0, \quad \lim_{j_+ \rightarrow 0} R = 0 && \text{if } U(1)_B \text{ broken} \end{aligned} \quad (3)$$

where the vanishing of R in the broken phase follows from the Goldstone nature of the qq_- boson in this case.




Figure 2. Susceptibility ratio R (left) and diquark condensate $\langle qq_+ \rangle$ (right) as functions of j_+ for various μ .

Figure 2 shows results for R and $\langle qq_+ \rangle$ from data taken on 12^4 , 16^4 and 20^4 lattices and then linearly extrapolated to the limit $L_t^{-1} = T \rightarrow 0$. For $\mu = 0$ the data appear to extrapolate smoothly to $R = 1$, $\langle qq_+ \rangle = 0$ as $j_+ \rightarrow 0$. For $\mu a = 1.0$ by contrast, which lies in the region of high baryon density and restored chiral symmetry, then if data with $j_+ a \leq 0.2$ are ignored the extrapolated values are $R \simeq 0$, $\langle qq_+ \rangle a^3 \simeq 0.28$, suggesting superfluidity. For intermediate μ the situation is not so clear cut, but the results of a quadratic extrapolation of $\langle qq_+ \rangle$ to $j_+ = 0$ are shown in Figure 1. For $0.4 \lesssim \mu a \lesssim 0.9$ the condensate scales as μ^2 , which is expected if diquark pairs within a distance Δ of a Fermi surface of radius μ participate in the condensate leading to $\langle qq_+ \rangle \sim \Delta \mu^2$. The value of $\langle qq_+ \rangle$ at $\mu a = 1.0$, together with an estimated $\Sigma \sim 350 \text{ MeV}$, is thus plausibly consistent with the gap values reported in [2]. The positive curvature contrasts with the behaviour of the superfluid condensate in Two Color QCD in which qq pairs are tightly bound and form a conventional Bose-Einstein condensate⁸.

Since our interpretation of broken $U(1)_B$ symmetry at high baryon density relies on discarding some data, the above conclusions are necessarily provisional. As spontaneous symmetry breaking does not occur in a finite

volume, it is natural to attribute the depletion of $\langle qq_+ \rangle$ at small j_+ to the effects of a finite L_s . Note however that the threshold $ja \sim 0.3$ required for behaviour characteristic of the thermodynamic limit seems anomalously large compared with that required to observe chiral symmetry breaking ($m_0 a \sim 0.002$) on the same volume, or indeed for thin film superfluidity ($ja \sim 0.1$) in NJL₂₊₁⁶. We suspect that finite volume effects in systems exhibiting a Fermi surface are unconventional, and indeed data on systems with varying L_s, L_t suggest the approach to the thermodynamic limit at small j is non-monotonic⁷. Solution of the gap equation on finite systems shows that $\Delta(L_s)$ oscillates before approaching its infinite volume limit for $L_s \gtrsim 8\text{fm}$, which translates to a 40^3 spatial lattice for our model⁹.

In future work we plan to study the quasiparticle spectrum which in principle permits a direct estimate of the superfluid gap Δ . It might also prove interesting to choose different bare parameters, since a small change in the cutoff might actually change the chiral symmetry restoring transition to first order⁴, making it easier to distinguish the properties of high- and low- μ phases. Finally, if our conclusion that NJL₃₊₁ is superfluid at high density persists, then it will be interesting to examine the stability of the condensate as either temperature or $\mu_{isospin}$ (which has the effect of separating u and d Fermi surfaces and hence inhibiting pairing in the isoscalar channel) is raised from zero.

References

1. K. Rajagopal and F. Wilczek, in *Handbook of QCD* ch. 35 ed. M. Shifman (World Scientific, Singapore 2001).
2. J. Berges and K. Rajagopal, *Nucl. Phys.* **B538**, 215 (1999).
3. S.J. Hands, *Nucl. Phys.* **A642**, 228 (1998).
4. S.P. Klevansky, *Rev. Mod. Phys.* **64**, 649 (1992).
5. S.J. Hands and S.E. Morrison, *Phys. Rev.* **D59**:116002 (1999).
6. S.J. Hands, B. Lucini and S.E. Morrison, *Phys. Rev. Lett.* **86**, 753 (2001); *Phys. Rev.* **D65**:036004 (2002).
7. S.J. Hands and D.N. Walters, [hep-lat/0209140](#).
8. J.B. Kogut, M.A. Stephanov, D. Toublan, J.J.M. Verbaarschot and A. Zhitnitsky, *Nucl. Phys.* **B582**, 477 (2000).
9. P. Amore, M.C. Birse, J.A. McGovern and N.R. Walet, *Phys. Rev.* **D65**:074005 (2002).